

REVIEW ARTICLE

Eco-Friendly Alternatives for Managing Plant Diseases: Lessons from *Ageratum conyzoides* -A Review

Saurav Anand¹, Rajeev Kumar^{2*}, Manjot Kaur³, Deepa Rani⁴

¹School of Bioengineering and Bioscience, Lovely Professional University, Phagwara-144 411, Punjab, India

²School of Agriculture, Lovely Professional University, , Phagwara, 144411, Punjab, India

³Department of Microbiology, DAV University, Jalandhar, Punjab, India- 144012

⁴Department of Pharmacognosy, M.M college of Pharmacy, Maharishi Markandeshwar (Deemed to be University), Mullana, Ambala, 133207

Received: 15th October, 2025; Revised: 30th October, 2025; Accepted: 12th November, 2025; Available Online: 25th December, 2025

ABSTRACT

Ageratum conyzoides, commonly known as billygoat weed or goatweed, poses a significant threat to agricultural ecosystems, negatively impacting crop productivity and biodiversity. Due to environmental concerns associated with conventional control methods, this comprehensive review critically examines the current state of biological control strategies for *A. conyzoides*. The study explores three major categories of biological control agents: insects such as the *Ageratum* gall fly (*Procecidochares utilis*) and the *Ageratum* stem-boring weevil (*Listronotus setosipennis*), which disrupt weed growth through gall induction and stem boring, respectively; fungal pathogens that induce diseases compromising weed vigor; and herbivorous livestock, particularly controlled grazing by goats and sheep, as a natural means of suppression. The review evaluates the efficacy of these agents while considering key environmental factors such as climate and soil characteristics. Special emphasis is placed on minimizing non-target effects through host specificity assessments and optimizing biocontrol implementation strategies. Additionally, challenges including the potential development of resistance in *A. conyzoides*, the necessity of integrating multiple control measures, and existing knowledge gaps are discussed. This study underscores the importance of continued research and the adoption of integrated pest management (IPM) approaches to achieve sustainable and ecologically sound weed control. In conclusion, this review provides valuable insights into eco-friendly weed management practices, contributing to sustainable agricultural production and biodiversity conservation.

Keywords: *Ageratum conyzoides*, biological control, weed management, insect pests, pathogens, sustainable agriculture, environmental conservation.

Micro Environer (2025); DOI: <https://doi.org/10.54458/mev.v5i02.04>

How to cite this article: Anand S, Kumar R, Kaur M, Rani D. Eco-Friendly Alternatives for Managing Plant Diseases: Lessons from *Ageratum conyzoides* -A Review. Micro Environer. 2025;5(2):18-22.

Source of support: Nil.

Conflict of interest: None

INTRODUCTION

Agriculture, a cornerstone of human civilization, continually confronts challenges from various pests and invasive species that threaten the productivity of cultivated lands and the equilibrium of ecosystems. *A. conyzoides*, colloquially known as billygoat weed or goatweed, has emerged as a formidable adversary in this intricate dance between human cultivation and the natural world. Native to tropical and subtropical regions, this tenacious weed possesses a remarkable ability to adapt and proliferate, casting its shadow over crops, pastures, and natural habitats.

As the global agricultural landscape grapples with the impact of conventional herbicides on the environment and con-

cerns about their sustainability, the exploration of alternative and eco-friendly control strategies becomes imperative. Singh, M. (2024). Bio- logical control, a time-honored approach, harnesses the inher- ent mechanisms of the natural world to manage pest popula- tions. In the context of *A. conyzoides*, a weed with a pervasive presence, the quest for effective biological control agents has gained momentum, driven by the need for sustainable solutions that harmonize with the delicate balance of ecosystems.

Ageratum conyzoides

A. conyzoides, a member of the Asteraceae family, stands as a testament to nature's resilience and adaptability. Ex-

hibiting a penchant for disturbed habitats, agricultural fields, and open spaces, this annual herbaceous plant has earned its reputation as a noxious weed. Its rapid growth, prolific seed production, and allelopathic properties contribute to its ability to outcompete native vegetation, compromising biodiversity and impacting crop yields. The invasiveness of *A. conyzoides* poses a significant threat to both agricultural productivity and the ecological integrity of diverse landscapes. Kumar, R *et al*, (2024).

Traditional methods of weed control, often reliant on chemical herbicides, come with a litany of concerns. Environmental contamination, the development of herbicide-resistant strains, and unintended harm to non-target organisms raise questions about the sustainability of these practices. Consequently, the imperative to explore alternative, environmentally benign strategies for *A. conyzoides* management has driven researchers, agriculturalists, and environmentalists towards the realm of biological control. Kumar, R *et al*, (2022)

Biological Control

Biological control, a branch of integrated pest management (IPM), harnesses the natural enemies of pests to limit their populations and mitigate their impact. In the case of *A. conyzoides*, biological control represents a promising avenue for sustainable weed management. By leveraging the interactions between organisms within ecosystems, this approach aims to restore a semblance of balance, allowing native flora to thrive while suppressing the invasive billygoat weed. Chen, S *et al*, (2025). The concept of biological control extends across various trophic levels, involving predators, parasitoids, pathogens, and herbivores. In the intricate dance of nature, certain organisms have evolved to exploit specific vulnerabilities of *A. conyzoides*, be it through feeding, parasitism, or inducing diseases. The focus on biological control aligns with broader trends in sustainable agriculture, emphasizing the need for holistic and environmentally friendly solutions.

Insects as Biological Control Agents

In the arsenal of biological control agents, insects emerge as key players in the endeavor to manage *A. conyzoides*. The *Ageratum* gall fly (Procecidochares utilis) and the *Ageratum* stem-boring weevil (Listronotus setosipennis) showcase the potential of insects to disrupt the weed's life cycle. The gall fly's intricate dance involves laying eggs on *A. conyzoides*, inducing gall formation that disrupts the weed's growth and reproduction. Simultaneously, the stem-boring weevil's larvae burrow into the stems, causing structural damage and reducing the vigor of the billygoat weed. Kato-Noguchi and kato, (2024).

These insects, acting as natural adversaries, exemplify the elegance of coevolution and the intricate mechanisms through which the natural world seeks equilibrium. The specificity of these insects to *A. conyzoides* minimizes the risk to non-target species, aligning with the principles of precision and sustainability inherent in biological control.

Pathogens

Fungal pathogens, another cohort of biological control agents, introduce a different dimension to the battle against

A. conyzoides. Certain fungi exhibit an aptitude for infecting the weed, causing diseases that compromise its growth and reproductive potential. This silent warfare beneath the soil surface not only weakens the individual plants but also curtails the spread of *A. conyzoides* by affecting its seed production.

The use of fungi as biocontrol agents is notable for its potential specificity to the target weed, minimizing the risk to non-target species. This approach aligns with the ecological principles of sustainability, offering a focused solution to the challenges posed by *A. conyzoides* in diverse ecosystems.

Livestock Grazing

Beyond the microscopic realm of insects and fungi, the integration of livestock, such as goats and sheep, introduces a macroscopic yet equally natural approach to *A. conyzoides* management. Grazing animals, with their voracious appetites for certain weeds, including *A. conyzoides*, offer a sustainable and economically viable solution. Controlled grazing not only reduces the biomass of billygoat weed but also contributes to nutrient cycling and promotes a diverse and resilient pasture ecosystem.

The use of livestock as biological control agents aligns with the principles of agroecology, where agriculture is viewed through an ecological lens, recognizing the interconnectedness of various components within the system. As these animals graze, they act as stewards of the land, participating in a natural symphony that echoes the principles of sustainable land management.

Challenges and Opportunities in Biological Control

While biological control holds immense promise in managing *A. conyzoides*, it is not without its challenges. The potential for the weed to develop resistance to biocontrol agents, concerns about unintended harm to non-target species, and the need for a nuanced understanding of the ecological dynamics within specific ecosystems are pivotal considerations. Addressing these challenges requires a multidisciplinary approach, blending ecological insights with advancements in entomology, plant pathology, and agronomy.

The integration of biological control into a broader weed management strategy becomes paramount in navigating these challenges. Recognizing that no single solution fits all scenarios, an approach that combines biological control with cultural, mechanical, and chemical methods, under the umbrella of integrated pest management (IPM), emerges as a comprehensive strategy. Li, Y *et al*, (2025).

Literature review

This literature review explores a selection of studies that investigate the antifungal, pesticidal, and allelopathic properties of *A. conyzoides* and its potential applications in sustainable agriculture.

Antifungal Activity

The study by Javed and Bashir (2012) investigates the antifungal activity of different extracts of *A. conyzoides* against *Fusarium solani*. Published in the African Journal

of Biotechnology, the research explores the potential of *A. conyzoides* as a natural agent for managing fungal pathogens, shedding light on its bioactive compounds.

Building upon this, Iqbal *et al.* (2004) contribute to the understanding of *A. conyzoides* fungistatic properties. Their work, published in *Phytoparasitica*, identifies a fungistatic chromene from *A. conyzoides*, showcasing the richness of bioactive compounds within the plant that may contribute to its antifungal properties.

In a more recent study, Banaras *et al.* (2021) perform bioassays guided fractionation of *A. conyzoides* extract to identify natural antifungal compounds against *Macrophomina phaseolina*. Published in the *International Journal of Agriculture and Biology*, this research highlights the potential of *A. conyzoides* in biocontrol against specific fungal pathogens.

Ndacnou *et al.* (2020) contribute to the phytochemical study of *A. conyzoides* and assess its anti-oomycete activity. Published in *Industrial Crops and Products*, their work expands our understanding of the plant's chemical composition and its relevance in managing oomycete infections.

Pest and Disease Management

Rioba and Stevenson (2017) explore the utilization of *A. conyzoides* for the management of pests and diseases by smallholder farmers. Published in *Industrial Crops and Products*, this study provides insights into the practical applications of *A. conyzoides* in real-world agricultural settings.

Chen *et al.* (2023) contribute to the field by screening and characterizing biocontrol bacteria isolated from *A. conyzoides* against *Colletotrichum fructicola*, a pathogen causing Chinese plum anthracnose. Their work, published in *Frontiers in Microbiology*, showcases the potential of *A. conyzoides* not only as a direct biocontrol agent but also as a source of beneficial microbes for integrated pest management.

Allelopathic Potential

Kong *et al.* (1999) and Kong *et al.* (2004) delve into the allelopathic potential of *A. conyzoides*. The studies, published



Figure 1 : Fungal strains sensitive to *Ageratum* extracts/oils

in the *Journal of Chemical Ecology* and *Allelopathy Journal*, respectively, explore the plant's ability to release allelopathic compounds that inhibit the growth of neighboring plants. This allelopathic potential holds promise for weed management in agroecosystems.

Multi-faceted Contributions

A comprehensive study by Chahal *et al.* (2021) examines *A. conyzoides* and its secondary metabolites in the management of different fungal pathogens. Published in *Molecules*, this research not only highlights its antifungal properties but also emphasizes the plant's potential against a spectrum of fungal adversaries.

Beyond Agriculture

A. conyzoides is not limited to agricultural applications alone. Paul *et al.* (2022) discuss its potential in turning waste into beneficial resources, implying implications in sustainable agriculture, the environment, and the biopharmaceuticals. Published in *Molecular Biotechnology*, this study broadens the scope of *A. conyzoides*' contributions.

CONCLUSION

The management of *Ageratum conyzoides* remains a critical challenge in sustainable agriculture, necessitating an integrative approach to mitigate its adverse effects on crop production and ecosystem stability. This review comprehensively elucidates the role of biocontrol strategies, including insect-mediated suppression, fungal antagonism, and livestock grazing, in effectively reducing the competitive dominance of this invasive weed. The utilization of *Procecidochares utilis* (*Ageratum* gall fly) and *Listronotus setosipennis* (*Ageratum* stem-boring weevil) demonstrates species-specific interactions that disrupt weed physiology, thereby limiting growth and reproductive potential. Concurrently, pathogenic fungi

Table 1: Antifungal Activities of Main Constituents Extracted from *Ageratum conyzoides*

Constituent	Antifungal Activity Against	Target Pathogen
Chromene	Fusarium solani	Fusarium solani
	Macrophomina phaseolina	Macrophomina phaseolina
Bioactive Compounds	Lasiodiplodia theobromae	Lasiodiplodia theobromae
	Lasiodiplodia pseudotheobromae	Lasiodiplodia pseudotheobromae
Essential Oil	Aspergillus spp.	Virulent Aspergillus spp.
	Phytophthora capsici	Phytophthora capsici
Allelochemicals	Various fungal pathogens	Various fungal pathogens

exert significant mycoherbicidal effects, causing structural degradation in plant tissues, ultimately leading to suppression of *A. conyzoides* populations. The allelopathic properties of *A. conyzoides*, attributed to bioactive secondary metabolites such as chromenes, flavonoids, and terpenoids, present both an ecological advantage and an agricultural constraint. While these compounds exhibit antifungal and insecticidal properties, their persistence in the soil matrix can influence native flora and soil microbiota, necessitating further investigation into their long-term ecological implications. Livestock grazing, particularly by goats and sheep, provides an ecologically sustainable weed suppression method, contributing to nutrient cycling and enhancing soil organic matter content. However, variable palatability and grazing preferences among livestock species require optimized grazing protocols to maximize efficacy. Integration of these biocontrol strategies within the framework of integrated weed management (IWM) offers a multifaceted approach to managing *A. conyzoides*. Synergistic combinations of biological, cultural, and chemical control methods, guided by ecological principles, can enhance long-term suppression while mitigating resistance development. Nevertheless, biocontrol implementation is constrained by environmental heterogeneity, host specificity of agents, and regulatory challenges in field application. Future research should emphasize molecular characterization of plant-microbe interactions, genomic insights into resistance mechanisms, and formulation of biopesticides derived from *A. conyzoides* extracts for targeted weed suppression. This review underscores the significance of biocontrol as an environmentally benign alternative to conventional herbicide-based weed management. Advancements in microbial consortia, gene-editing technologies for pest resistance, and precision agriculture tools hold promise for refining biocontrol efficacy. The holistic adoption of sustainable weed management strategies will ensure ecological balance, enhance crop resilience, and contribute to long-term agricultural productivity.

REFERENCES

- Asman, A., Cahyani, A., Nufus, A., Rosmana, A., Fakhruddin, A., & Natsir, N. (2021). Antifungal activity of extracts of *Melia azedarach* and *Ageratum conyzoides* against *Lasiodiplodia pseudotheobromae* through in vitro test. In IOP Conference Series: Earth and Environmental Science (pp. 012007).
- Asman, A., Rosmana, A., Sjam, S., Fakhruddin, A., Natsir, N., & others (2021). Antifungal activity of crude extracts of *Ageratum conyzoides* and *Chromolaena odorata* for management of *Lasiodiplodia theobromae* and *Lasiodiplodia pseudotheobromae* through in vitro evaluation. In IOP Conference Series: Earth and Environmental Science (pp. 012008).
- Banaras, S., Javaid, A., & Khan, I. (2021). Bioassays guided fractionation of *Ageratum conyzoides* extract for the identification of natural antifungal compounds against *Macrophomina phaseolina*. International Journal of Agriculture and Biology, 25(4), 761–767.
- Batish, D., Kaur, S., Singh, H., & Kohli, R. (2009). Nature of interference potential of leaf debris of *Ageratum conyzoides*. Plant Growth Regulation, 57, 137–144.
- Borah, R., Borah, J., & Islam, S. (2019). Biocontrol of five invasive weeds of Meghalaya-A case study. Journal of Biological Control, 33(2).
- Bouda, H., Tapondjou, L., Fontem, D., & Gumedzoe, M. (2001). Effect of essential oils from leaves of *Ageratum conyzoides*, *Lantana camara* and *Chromolaena odorata* on the mortality of *Sitophilus zeamais* (Coleoptera, Curculionidae). Journal of Stored Products Research, 37(2), 103–109.
- Chahal, R., Nanda, A., Akkol, E., Sobarzo-Sanchez, E., Arya, A., Kaushik, D., Dutt, R., Bhardwaj, R., Rahman, M., & Mittal, V. (2021). *Ageratum conyzoides* L. and its secondary metabolites in the management of different fungal pathogens. Molecules, 26(10), 2933.
- Chawla, R., Rajvir, S., Sunder, S., & others (2013). Isolates and herbicidal activity of fractions and extract of stems of *Ageratum conyzoides*. Annals of Agri Bio Research, 18(1), 82–86.
- Chen, X., Zhang, M., Tang, L., Huang, S., Guo, T., & Li, Q. (2023). Screening and characterization of biocontrol bacteria isolated from *Ageratum conyzoides* against *Collectotrichum fructicola* causing Chinese plum (*Prunus salicina* Lindl.) anthracnose. Frontiers in Microbiology, 14.
- Devi, O., Dutta, B., & Choudhury, P. (2013). Allelopathy effect of aqueous extract of *Clerodendrum viscosum*, *Ageratum conyzoides* and *Parthenium hysterophorus* on the seed germination and seedling vigour of Chickpea seeds (*Cicer arietinum* L.) in vitro. Journal of Applied and Natural Science, 5(1), 37–40.
- Fagoonee, I., Umrit, G., & others (1980). Biology of *Dysdercus cingulatus* Sign. and its control by *Ageratum conyzoides*. Revue Agricole et sucriere de l'Île Maurice, 59(3), 122–128.
- Iqbal, M., Jayasinghe, U., Herath, H., Wijesekara, K., & Fujimoto, Y. (2004). A fungistatic chromene from *Ageratum conyzoides*. Phytoparasitica, 32, 119–126.
- Javed, S., & Bashir, U. (2012). Antifungal activity of different extracts of *Ageratum conyzoides* for the management of *Fusarium solani*. African Journal of Biotechnology, 11(49), 11022–11029.
- Kohli, R., Batish, D., Singh, H., & Dogra, K. (2006). Status, invasiveness and environmental threats of three tropical American invasive weeds (*Parthenium hysterophorus* L., *Ageratum conyzoides* L., *Lantana camara* L.) in India. Biological Invasions, 8, 1501–1510.
- Kong, C. (2010). Ecological pest management and control by using allelopathic weeds (*Ageratum conyzoides*, *Ambrosia trifida*, and *Lantana camara*) and their allelochemicals in China. Weed Biology and Management, 10(2), 73–80.
- Kong, C., Hu, F., & Xu, X. (2002). Allelopathic potential and chemical constituents of volatiles from *Ageratum conyzoides* under stress. Journal of Chemical Ecology, 28, 1173–1182.
- Kong, C., Hu, F., Xu, X., Liang, W., & Zhang, C. (2004). Allelopathic plants. *Ageratum conyzoides* L. Allelopathy Journal, 14, 1–12.
- Kong, C., Hu, F., Liang, W., Peng, W., Jiang, Y., & others (2004). Allelopathic potential of *Ageratum conyzoides* at various growth stages in different habitats. Allelopathy Journal, 13(2), 233–240.
- Kong, C., Liang, W., Hu, F., Xu, X., Wang, P., Jiang, Y., & Xing, B. (2004). Allelochemicals and their transformations in the *Ageratum conyzoides* intercropped citrus orchard soils. Plant and Soil, 264, 149–157.
- Kumar, R., Singh, J. P., Rana, M., & Srivastava, S. (2024).

Antifungal potential of allelopathic plant extracts of *Ageratum conyzoides* L. and *Parthenium hysterophorus* L. on Phytophthora blight of *Capsicum annuum* L. *Allelopathy Journal*, 63(2).

21. Ming, L. (1999). *Ageratum conyzoides*: A tropical source of medicinal and agricultural products. *Perspectives on New Crops and New Uses*, 1988, 469–473.

22. Mukhtar, I., Khokhar, I., & Mushtaq, S. (2013). New record of *Podosphaera xanthii* on *Ageratum conyzoides* L. in Pakistan. *Pakistan Journal of Weed Science Research*, 19(4).

23. Ndacnou, M., Pantaleon, A., Tchinda, J.b., Mangapche, E., Keumedjio, F., & Boyoguemo, D. (2020). Phytochemical study and anti-oomycete activity of *Ageratum conyzoides* Linnaeus. *Industrial Crops and Products*, 153, 112589.

24. Nogueira, J., Gonçalez, E., Galleti, S., Facanali, R., Marques, M., & Felício, J. (2010). *Ageratum conyzoides* essential oil as aflatoxin suppressor of *Aspergillus flavus*. *International Journal of Food Microbiology*, 137(1), 55–60.

25. Okunade, A. (2002). *Ageratum conyzoides* L.(Asteraceae). *Fitoterapia*, 73(1), 1–16.

26. Paul, S., Datta, B., Ratnaparkhe, M., & Dholakia, B. (2022). Turning waste into beneficial resource: Implication of *Ageratum conyzoides* L. in sustainable agriculture, environment and biopharma sectors. *Molecular Biotechnology*, 64(3), 221–244.

27. Pavaraj, M., Karthikairaj, K., & Rajan, M. (2010). Effect of leaf extract of *Ageratum conyzoides* on the biochemical profile of blackgram *Vigna mungo* infected by root-knot nematode, *Meloidogyne incognita*. *Journal of Biopesticides*, 3(Special Issue), 313.